



Genome Sequence of "Candidatus Methanomethylophilus alvus" Mx1201, a Methanogenic Archaeon from the Human Gut Belonging to a Seventh Order of Methanogens

Guillaume Borrel, Hugh Mb Harris, William Tottey, A. Mihajlovski, Nicolas Parisot, Eric Peyretailade, P. Peyret, Simonetta Gribaldo, Paul W. O'Toole, Jean François Brugère

► To cite this version:

Guillaume Borrel, Hugh Mb Harris, William Tottey, A. Mihajlovski, Nicolas Parisot, et al.. Genome Sequence of "Candidatus Methanomethylophilus alvus" Mx1201, a Methanogenic Archaeon from the Human Gut Belonging to a Seventh Order of Methanogens. *Journal of Bacteriology*, 2012, 194 (24), pp.6944 - 6945. 10.1128/JB.01867-12 . hal-01612754

HAL Id: hal-01612754

<https://hal.science/hal-01612754>

Submitted on 8 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genome Sequence of "Candidatus Methanomethylophilus alvus" Mx1201, a Methanogenic Archaeon from the Human Gut Belonging to a Seventh Order of Methanogens

Guillaume Borrel, Hugh M. B. Harris, William Tottey, Agnès Mihajlovski, Nicolas Parisot, Eric Peyretailade, Pierre Peyret, Simonetta Gribaldo, Paul W. O'Toole and Jean-François Brugère
J. Bacteriol. 2012, 194(24):6944. DOI: 10.1128/JB.01867-12.

Updated information and services can be found at:
<http://jb.asm.org/content/194/24/6944>

	<i>These include:</i>
REFERENCES	This article cites 17 articles, 6 of which can be accessed free at: http://jb.asm.org/content/194/24/6944#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Genome Sequence of “*Candidatus Methanomethylophilus alvus*” Mx1201, a Methanogenic Archaeon from the Human Gut Belonging to a Seventh Order of Methanogens

Guillaume Borrel,^a Hugh M. B. Harris,^b William Tottey,^a Agnès Mihajlovski,^a Nicolas Parisot,^a Eric Peyretailade,^a Pierre Peyret,^a Simonetta Gribaldo,^c Paul W. O’Toole,^b and Jean-François Brügère^a

EA-4678 CIDAM, Conception Ingénierie et Développement de l’Aliment et du Médicament, Clermont Université, Université d’Auvergne, Clermont-Ferrand, France^a; Department of Microbiology and Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland^b; and Unité de Biologie Moléculaire du Gène chez les Extrémophiles, Institut Pasteur, Paris, France^c

We report the draft genome sequence of “*Candidatus Methanomethylophilus alvus*” Mx1201, a methanogen present in the human gut. It was enriched from human feces under anaerobic conditions with methanol as the substrate. Its circular genome, of around 1.7 Mb, contains genes needed for methylotrophic methanogenesis from methanol and tri-, di-, and monomethylamine.

Methanogenic archaea play a crucial role in the functioning of digestive ecosystems through the consumption of the hydrogen generated during anaerobic fermentations (3). The first archaeal species associated to the human gut, *Methanobrevibacter smithii* and *Methanosphaera stadtmanae* (order *Methanobacteriales*), were cultured and described 30 years ago (12, 13). During the last decade, the diversity of the human-associated archaea was progressively enlarged by the detection of 16S rRNA genes affiliated to the *Crenarchaeota* (15), to the *Halobacteriales* (14), and to a *Thermoplasmatales*-related lineage (10, 11, 16). A species of the latter lineage, *Methanomassiliicoccus luminyensis*, was recently isolated and described as a methanogen growing on methanol and hydrogen only (5). More recently, its genome was sequenced (6).

To allow genomic characterization of novel methanogens, we performed enrichment cultures targeting methanogens in feces of elderly people (68 to 93 years old), selecting for the presence of a *Thermoplasmatales*-related lineage. Enrichment cultures carried out with methanol as the substrate led to the progressive dominance of “*Candidatus Methanomethylophilus alvus*” over the microbial community. This archaeon is distantly related to its closest related species, *M. luminyensis* and *Aciduliprofundum boonei*, with 87% and 80% 16S rRNA gene sequence identity, respectively. Sequences of 16S rRNA genes of uncultured archaea with more than 97% identity to “*Ca. Methanomethylophilus alvus*” have previously been recurrently retrieved in the digestive tract of various animals, suggesting that these archaea are specifically associated to these environments. Those sequences form a portion of the previously described rumen cluster C (8).

A 3-kb mate-paired library was constructed and sequenced from a quarter plate of a 454 GS FLX Titanium run (Macrogen, Republic of Korea). A total of 347,052 reads corresponding to 158 Mb were obtained. The reads were assembled with a GS De Novo assembler (version 2.6), first in 15 contigs (average depth coverage of 36.9-fold) and then in a unique scaffold of 1,664,410 bp. The gaps between the contigs were partially closed using GapFiller (2) and by sequencing of PCR products (4 remaining gaps). Around 1,800 bp was lacking between the two scaffold termini, as shown by PCR amplification. The open reading frames were predicted with Glimmer 3 (4) and annotated using RAST (1).

The genome of “*Ca. Methanomethylophilus alvus*” has a G+C content of 55.6%, 44 usual tRNA genes, a single copy of each of the

23S and 16S rRNA genes, and 2 noncontiguous copies of 5S rRNA genes that were distant in the genome from the 23S and 16S rRNA genes. A total of 1,785 protein coding sequences were predicted. The identification of the 56 ribosomal proteins ubiquitous in archaeal genomes (17) confirms the near completeness of the genome.

Two confidently annotated CRISPR regions (and two candidates) were identified using CRISPRFinder (7), in close association with *cas* genes (encoding CRISPR-associated proteins).

The presence of genes involved in methanogenesis, among them the highly conserved *mcrA* gene, supports the experimental observations that “*Ca. Methanomethylophilus alvus*” is a methanogen. Genes needed for methylotrophic methanogenesis from methanol and tri-, di-, and monomethylamine compounds were also identified. The amber codon in methyltransferases genes (*mtmB*, *mtbB* and *mttB*), together with genes involved in pyrrolysine (Pyl) biosynthesis (the 22nd amino acid carried by a tRNA^{Pyl} with a CUA anticodon), suggests the presence of Pyl in these proteins (9). “*Ca. Methanomethylophilus alvus*” would not only be able to reduce methanol with H₂ for methanogenesis but could probably also reduce other methylated compounds. To our knowledge, methylamines have not yet been considered substrates for methanogenesis, *in extenso* as valuable electron acceptors, in the human gut.

Nucleotide sequence accession numbers. The draft genome sequence of “*Ca. Methanomethylophilus alvus*” Mx1201 has been deposited at DDBJ/EMBL/GenBank under the accession no. [AMSE000000000](#). The version described in this paper is the first version, [AMSE010000000](#).

ACKNOWLEDGMENTS

We are grateful to A. Mansoor, P. Denozi, and their team of the “Centre Hospitalier Paul Ardière” in Issoire, France, for their valuable help in collecting fecal samples.

Received 28 September 2012 Accepted 1 October 2012

Address correspondence to Jean-François Brügère, j-francois.Brugere@udamail.fr.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01867-12

P.W.O. was supported by Science Foundation Ireland through a Principal Investigator award and by a Centre for Science, Engineering, and Technology award to the Alimentary Pharmabiotic Centre, Cork, Ireland. The authors have no conflicts of interests to declare.

REFERENCES

1. Aziz R, et al. 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
2. Boetzer M, Pirovano W. 2012. Toward almost closed genomes with GapFiller. *Genome Biol.* 13:R56.
3. Carbonero F, Benefiel AC, Gaskins HR. 2012. Contributions of the microbial hydrogen economy to colonic homeostasis. *Nat. Rev. Gastroenterol.* 9:504–518.
4. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23:673–679.
5. Dridi B, Fardeau ML, Ollivier B, Raoult D, Drancourt M. 2012. *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. *Int. J. Syst. Evol. Microbiol.* 62:1902–1907.
6. Gorlas A, Robert C, Gimenez G, Drancourt M, Raoult D. 2012. Complete genome sequence of *Methanomassiliicoccus luminyensis*, the largest genome of a human-associated Archaea species. *J. Bacteriol.* 194:4745–4745.
7. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a Web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res.* 35:W52–W57.
8. Janssen PH, Kirs M. 2008. Structure of the archaeal community of the rumen. *Appl. Environ. Microbiol.* 74:3619–3625.
9. Krzycki JA. 2005. The direct genetic encoding of pyrrolysine. *Curr. Opin. Microbiol.* 8:706–712.
10. Mihajlovski A, Alric M, Brugere JF. 2008. A putative new order of methanogenic Archaea inhabiting the human gut, as revealed by molecular analyses of the *mcrA* gene. *Res. Microbiol.* 159:516–521.
11. Mihajlovski A, Dore J, Levenez F, Alric M, Brugere JF. 2010. Molecular evaluation of the human gut methanogenic archaeal microbiota reveals an age-associated increase of the diversity. *Environ. Microbiol. Rep.* 2:272–280.
12. Miller TL, Wolin M, de Macario EC, Macario A. 1982. Isolation of *Methanobrevibacter smithii* from human feces. *Appl. Environ. Microbiol.* 43:227–232.
13. Miller TL, Wolin MJ. 1985. *Methanosphaera stadtmaniae* gen. nov., sp. nov.: a species that forms methane by reducing methanol with hydrogen. *Arch. Microbiol.* 141:116–122.
14. Oxley A, et al. 2010. Halophilic archaea in the human intestinal mucosa. *Environ. Microbiol.* 12:2398–2410.
15. Rieu-Lesme F, Delbès C, Sollelis L. 2005. Recovery of partial 16S rDNA sequences suggests the presence of Crenarchaeota in the human digestive ecosystem. *Curr. Microbiol.* 51:317–321.
16. Scanlan P, Shanahan F, Marchesi J. 2008. Human methanogen diversity and incidence in healthy and diseased colonic groups using *mcrA* gene analysis. *BMC Microbiol.* 8:79.
17. Yutin N, Puigbò P, Koonin EV, Wolf YI. 2012. Phylogenomics of prokaryotic ribosomal proteins. *PLoS One* 7:e36972. doi:10.1371/journal.pone.0036972.